

IDAHO PREVENTIVE CARE RECOMMENDATIONS ADULT- AGES 19 AND ABOVE

REFERENCES

Key references used to develop the guidelines include:

- Clinicians Handbook of Preventive Services, 2nd Edition, 1998
- Guide to Clinical Preventive Services, Report of the U. S. Preventive Services Task Force, 2nd Edition, 1996 (USPSTF) and 3rd Edition, September 2002
- USPSTF Updates from the AHRQ Preventive Services Webpage at <https://www.uspreventiveservicestaskforce.org/Page/Name/recommendations>
- National Guideline Clearinghouse from the AHRQ Preventive Services Webpage at <https://www.guideline.gov/> (Funding for upkeep of this site is currently not available).
- Guidelines for Women's Health Care, The American College of Obstetricians and Gynecologists, 1996
- [Women's Preventive Services Initiative \(ACOG\)](#)
- National Heart Lung and Blood Institute, NIH
 - [The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure \(JNC 7\)](#)
 - Identification, Evaluation and Treatment of Overweight and Obesity in Adults, October 2000, NIH Publication Number 00-4084
- ACC/AHA
 - 2013 Guideline on the Assessment of Cardiovascular Risk
 - [2017 Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults](#)
- American Diabetes Association
 - Screening for Diabetes, Diabetes Care 2002 Jan; 25(suppl1):s21-s24
- American Medical Association
 - [Assessment and Management of Adult Obesity](#): A Primer for Physicians, 2003
- National Osteoporosis Foundation
 - [Clinician's Guide to Prevention and Treatment of Osteoporosis, 2014](#)

- Ito K, et al. Possibility of re-screening intervals of more than one year in men with PSA levels of 4.0 ng/ml or less. Prostate 2003 Sep 15;57(1):8-13.
- The Center for Disease Control and Prevention
 - [Morbidity and Mortality Weekly Publications](#)
 - [National Center for Immunization and Respiratory Diseases](#)