



Management of Geriatric Hip Fracture

- ASA American Society of Anesthesiologists. ASA Physical Classification System. Available at: <https://www.asahq.org/resources/clinical-information/asa-physical-status-classification-system>. Updated October 15, 2014. Accessed March 1, 2017.
- BRA Brauer CA, Coca-Perrillon M, Cutler DM, Rosen AB. Incidence and mortality of hip fractures in the United States. *JAMA*. 2009;302:1573-1579.
- BUT Butler M, Forte M, Kane RL, et al. *Treatment of Common Hip Fractures*. National Institutes of Medicine. Agency for Healthcare Research and Quality. AHRQ Publication 09-E013. Available at: https://www.ncbi.nlm.nih.gov/books/NBK32591/pdf/Bookshelf_NBK32591.pdf. Accessed January 18, 2017.
- CMS1 Department of Health and Human Services, Centers for Medicare and Medicaid Services. Chapter 1: Inpatient hospital services covered under part A. In: *Medicare Benefit Policy Manual* (Rev. 189), 2014: 110.2-110.3. Available at <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c01.pdf>. Accessed December 20, 2016.
- CMS2 Center for Medicare and Medicaid Services, Department of Health and Human Services. *Skilled Nursing Facility (SNF) Billing Reference*. 2015. Available at <https://www.cms.gov/Outreach-and-Education/Medicare-Learning-Network-MLN/MLNProducts/downloads/snfspellillnesschrt.pdf>. Accessed March 18, 2016
- DEJ de Jong MR, Van der Elst M, Hartholt KA. Drug-related falls in older patients: Implicated drugs, consequences, and possible prevention strategies. *Ther Adv Drug Saf*. 2013;4(4):147-154.
- FTO Ftouh S, Morga A, Swift C, et al. Management of hip fracture in adults: Summary of NICE guidance. *BMJ*. 2011;342:d3304.
- HAY Haybach PJ. Ototoxicity. Vestibular Disorders Association website. <http://vestibular.org/ototoxicity>. Published December 28, 2015. Accessed January 18, 2017.
- HOR Horlocker TT, Wedel DJ, Rowlingson JC, et al. Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: ASRA practice advisory (3rd ed). *Reg Anesth Pain Med*. 2010;35(1):64-101.
- KUC Kucher N, Koo S, Quiroz R, et al. Electronic alerts to prevent venous thromboembolism among hospitalized patients. *N Engl J Med*. 2005;352(10):969-977.
- MAL Malviya A, Martin K, Harper I, et al. Enhanced recovery program for hip and knee replacement reduces death rate. *Acta Orthop*. 2011;82(5):577-558.
- MOH Mohanty S, Rosenthal RA, Russell MM, Neuman MD, Ko CY, Esnaola NF. Optimal perioperative management of the geriatric patient: A best practice guideline from the American College of Surgeons NSQIP and the American Geriatrics Society. *J Am Coll Surg*. 2016;222(5):930-947.
- NIK Nikkel LE, Fox EJ, Black KP, Davis C, Andersen L, Hollenbeak CS. Impact of comorbidities on hospitalization costs following hip fracture. *J Bone Joint Surg Am* 2012;94(1):9-17.
- POR Portegijs E, Edgren J, Salpakoski A, et al. Balance confidence was associated with mobility and balance performance in older people with fall-related hip fracture: A cross-sectional study. *Arch Phys Med Rehabil*. 2012;93(12):2340-2346.
- ROB Roberts KC, Brox TW, Jevsevar DS, Sevarino K. Management of hip fracture in the elderly. *J Am Acad Orthop Surg*. 2015;23:131-137.
- SIH Sihvonon S, Kulmala J, Kallinen M, Alén M, Kiviranta I, Sipila S. Postural balance and self-reported balance confidence in older adults with a hip fracture history. *Gerontology*. 2009;55(6):630-636.
- SMI Smith CM, Cotter V. Age-related changes in health. In: Boltz M, Capezuti E, Fulmer T, Zwicker D, editor(s). *Evidence-based geriatric nursing protocols for best practice*. 4th ed. New York (NY): Springer Publishing Company; 2012:23-47.